

GEOTHERMAL TRAINING PROGRAMME Orkustofnun, Grensasvegur 9, IS-108 Reykjavik, Iceland

APPENDICES TO THE REPORT: RESISTIVITY STRUCTURE OF PAKA GEOTHERMAL PROSPECT IN KENYA


by

Raymond M. Mwakirani

Geothermal Development Company Ltd P.O BOX 17700-20100 Nakuru KENYA rmwakirani@gdc.co.ke


United Nations University Geothermal Training Programme Reykjavík, Iceland Published in 2011 These are appendices to the report "Resistivity Structure of Paka geothermal prospect in Kenya" by Raymond M. Mwakirani at the UNU Geothermal Training Programme in 2011. Appendix I shows the TEM profiles from the Paka prospect and Occam inversion of the resistivity data. Appendix II shows processed data for all the MT soundings with various MT parameters extracted from the EDI files. Appendix III shows the TEM and MT resistivity data from the Paka prospect and 1-D joint inversion of the MT and TEM profiles and the corresponding model curves. Appendix IV contains additional iso-resistivity maps from the Paka prospect to ones shown in the main report. Appendix V shows additional resistivity cross-sections from the Paka prospect and their locations.


APPENDIX I: TEM profiles from the Paka prospect and Occam inversion of the resistivity data





 $t^{1/2}$ [µs]; Depth [m]


 $t^{1/2}$ [µs]; Depth [m]








































APPENDIX II: Processed data for all the MT soundings with various MT parameters extracted from the EDI file namely the apparent resistivity, and phase in the xy and yx directions plus the one calculated from determinant of the rotationally invariant tensor, electrical strike direction (Zstrike), coherency (measure of the data quality), skew (grey dots) is a measure of 2 dimensionality, tipper magnitude and ellipticity (open circles).



































































40


































































































































































































APPENDIX III: TEM and MT resistivity data from the Paka prospect and 1-D joint inversion of the MT and TEM profiles and the corresponding model curves




PAK33 o




PAK34 o




PAK36_o




PAK38 o




PAK45 o




PAK48_o




PAK53_o




PAK55R o




PAK57 o




PAK61 o




PAK62_o




PAK67_o









PAK70 o




PAK76_o




PAK77 o









PAK80 o




PAK82_o




PAK85 o









PAK88 o




PAK99_o




PAK105B o




PakMT01_o




PakMT03_o




PakMT05_o




PakMT09 o




PakMT11_o




PakMT12_o




PakMT13_o




PakMT14_o




PakMT15_o




PakMT55R_o




PakMT62 o




PakMT67_o




PakMT68_o




PakMT77 o




PakMT82_o




PakMT84_o




PakMT88_o





PakMT99 o





APPENDIX IV: Additional iso-resistivity maps from the Paka prospect














APPENDIX V: Additional resistivity cross-sections from the Paka prospect and their locations

132




FIGURE 12: Location of TEM soundings (green dots) and MT soundings (red dots) considered in this survey and black lines the location of the cross-sections used in interpreting the resistivity structure of the survey area




FIGURE 13: Resistivity cross-section E_W_3 in the Paka prospect based on 1D joint inversion of TEM and determinant of MT data




FIGURE 14: Resistivity cross-section E_W_4 in the Paka prospect based on 1D joint inversion of TEM and determinant of MT data




FIGURE 15: Resistivity cross-section *N_S_I* in the Paka prospect based on 1D joint inversion of TEM and determinant of MT data